INDIAN JOURNAL OF POPULATION AND DEVELOPMENT ISSN: 2583-4827; Volume 4(2), July 2024: 183-199

Prevalence and Socioeconomic Determinants of Hypertension among Women in India: A Crosssectional Study from the Nationally Representative Data

Mriganka Dolui Sanjit Sarkar Sameer Kumar Jena

Abstract

Hypertension is a global health concern and is emerging as a prominent noncommunicable disease in India. It has substantial implications for the health of women. This paper analyses demographic and socioeconomic determinants of hypertension among Indian women aged 15-49 years based on the data from the fifth round of the National Family Health Survey. The analysis reveals that the prevalence of hypertension increases with age which aligns with the global trend. Education plays a pivotal role, as women with lower educational levels are at higher risk of hypertension than women with higher levels of education. The prevalence of hypertension has also been found to be relatively higher in women with high standard of living and women residing in the rural areas. Obesity, diabetes, and smoking are found to be strong predictors of hypertension. The geospatial analysis reveals regional disparities, with higher prevalence in north-eastern and northern districts of the region. There is a need of adopting multifaceted approaches to addressing the problem of hypertension in India women. The findings of the study may serve as a vital resource for policymakers and public health practitioners for addressing the growing epidemic of hypertension in the country.

Introduction

According to the World Health Organization, there are five important risk factors for non-communicable diseases (NCDs), one of which is hypertension (Ezzati et al, 2002). Hypertension plays a substantial role in elevating the disability-adjusted life years (DALYs) related to cardiovascular diseases when compared to other metabolic risk factors such as elevated total cholesterol, increased fasting blood sugar level, and a high body mass index (BMI) (Ezzati et al, 2002). Hypertension is a severe and chronic medical condition that promotes the risk of mortality related to cardiovascular and kidney diseases. This condition arises when the pressure within blood vessels becomes excessively elevated (Abarca-Gómez et al, 2017). The prevalence of hypertension is on the rise in recent years, and it has emerged as a particularly concerning issue among women globally and in India (Anchala et

al, 2014). Approximately four out of every five people with hypertension do not receive adequate treatment of hypertension. It is, however, estimated that if countries can scale up the coverage, 76 million deaths may be averted between 2023 and 2050 (WHO, 2023). This is a matter of great importance, as health and well-being of women not only affect their own lives but have far-reaching implications for families and communities. The health consequences of hypertension can be compounded by other factors that increase the odds of heart attack, stroke, and kidney failure. These factors include tobacco use, unhealthy diet, harmful use of alcohol, lack of physical activity, exposure to persistent stress as well as obesity, high cholesterol, and diabetes mellitus (WHO, 2015).

With its diverse population and complex healthcare landscape, India presents a unique context for studying the prevalence and determinants of hypertension among women. Hypertension is a leading non-communicable disease in India (Dolui et al, 2023; Gupta, 2004) and the prevalence of hypertension in women aged 15-49 years is increasing (Bhimarasetty et al, 2022). Available studies substantiate the link between hypertension and BMI and family history of non-communicable diseases (Daniel et al, 2022). Among young adults aged 20-39 years, one in every nine, and in the general adult population, over 25 per cent have been found to be suffering from hypertension while one third of the young adults were found to be pre-hypertensive (Geevar et al, 2022). Among women aged 15 years and older, approximately 21 per cent have been diagnosed with hypertension, while 39 per cent are in the pre-hypertensive category (Government of India, 2021).

India has recently intensified its commitment to tackling hypertension on a population scale by introducing a population-based screening programme for hypertension, diabetes, and cancers along with efforts to reinforce primary and secondary healthcare facilities (Government of India, 2022). The management of hypertension in the community requires regular blood pressure evaluation to allow for the earliest possible introduction of secondary prevention measures (Saha et al. 2008). Screening of for psychological distress in high-risk hypertensive patients, particularly focusing on those who struggle to attain adequate blood pressure control or are apprehensive about the potential complications associated with hypertension. It is estimated that 25 per cent of hypertensive patients exhibited signs of psychological distress, including symptoms associated with depression, anxiety, or stress. (Loke and Ching, 2022). Usual habits of an individual also play an important role as smoking and using smokeless tobacco and alcohol consumption have been found to be the risk factors of hypertension in women (Mishra et al, 2022). Socioeconomic and lifestyle factors are also found to be important reasons for the difference in the prevalence of hypertension in urban and rural India (Boro and Banerjee, 2022a).

Data and Methods

The analysis is based on the data available from the Round 5 of the National Family Health Survey (NFHS-5) which was conducted during the period 2019-2021) by the International Institute for Population Sciences (IIPS), Mumbai under the aegis of the Government of India, Ministry of Health and Family Welfare. The survey covered the entire country and covered 6,36,699 households, 1,01,839 men aged 15-54 years, and 7,24,115 eligible women aged 15-49 years. Estimates of the prevalence of hypertension are available from the survey for the country, for states/Union Territories and for 707 districts, with reference to the status as of March 31st, 2017. Details of NFHS-5 including the sampling frame, survey design, and data collection process are discussed elsewhere (Government of India, 2022). The present analysis is limited to the data on systolic and diastolic blood pressure measured from 7,24,115 women aged 15-49 years covered during the survey. Based on the level of systolic and diastolic blood pressure, a woman was classified as hypertensive if at least one of four conditions were satisfied: 1) woman reported high blood pressure on two or more occasions by a doctor or other health professional, 2) woman taking prescribed medicine to lower blood pressure at the time of the survey, 3) woman had an average systolic blood pressure (SBP) \geq 140 mmHg, 4) woman had an average diastolic blood pressure (DBP) \geq 90 mmHg at the time of the survey. The blood pressure of the women was measured three times with an interval of five minutes using Omron Blood Pressure Monitor during the survey and the average of the three measurements are taken for the purpose of classification.

The outcome variable of the present analysis is a dichotomous variable which takes value 1 if a woman is hypertensive and 0 otherwise. On the other hand, a set of explanatory variables including age, education, religion, caste, standard of living, residence, Body Mass Index (BMI), whether woman is diabetic or not, type of cooking fuel used in the house, smoking behaviour, experience of passive Smoking, and use of tobacco. Bivariate and multivariate statistical methods have been used to explore the determinants of hypertension. Descriptive statistics are used to understand the variation in the prevalence of hypertension by background characteristics. Logistic regression analysis has been used to assess how different explanatory variables influence the prevalence of hypertension. The multicollinearity test using variance inflation factors (VIF) is applied to root out multicollinearity among explanatory variables. Finally, geospatial patterning of the prevalence of hypertension was carried using district level data. All statistical analysis and spatial mapping exercise was performed using STATA 17.0 and ArcGIS 10.8.2 software.

Results

Table 1 provides background characteristics of women aged 15-49 years covered during the 1920-2021 round of NFHS. A large proportion of women was in the age group 15-29 years. More than half of the women had at least higher secondary level education. More than three-fourth were Hindu while around 43 per cent belonged to Other Backward Classes. More than two-third were residing in rural areas. The prevalence of obesity, as measured by BMI, was around 24 per cent while around 18 per cent of women had low BMI. The prevalence of diabetes was 2.6 per cent, around 4 per cent were smokers, and around 3 per cent were consuming tobacco. Around 48 per cent women experienced passive smoking. Around 44 per cent were exposed to polluting cooking fuel.

Table 2 presents the prevalence of hypertension among women by their background characteristics. The overall prevalence of hypertension was around 14 per cent but it was higher in women aged 45-49 years. The prevalence of hypertension was higher in women with no education, women belonging to other religions and Other Social Classes.

The prevalence of hypertension was higher in urban obese women than in rural obese women, and in smokers and tobacco chewers. There the prevalence of hypertension increases with the standard of living of the woman, The prevalence of hypertension was the lowest among women with the poorest standard of living.

Background Characteristics	Women surveyed		
	Per cent	Number	
Age			
15-29	49.60	359559	
30-44	38.63	280497	
45-49	11.77	84059	
Education			
No Education	22.43	167304	
Primary	11.73	84983	
Secondary	50.18	370012	
Higher	15.65	101816	
Religion			
Hindu	81.36	546007	
Muslim	13.48	90729	
Others	5.16	87379	
Caste			
Scheduled Caste	21.89	139957	
Scheduled Tribe	9.29	135239	
Other Backward Classes (OBC)	42.92	276881	
Others	25.91	172038	
Wealth Index			
Poorest	18.50	149844	
Poorer	20.00	160340	
Middle	20.52	151505	
Richer	20.81	139607	
Richest	20.17	122819	
Region			
North	14.11	147615	
Central	31.04	184017	
East	16.61	104342	
Northeast	3.69	103433	
West	13.94	69811	
South	20.60	114897	
Residence			
Urban	32.49	179535	
Rural	67.51	544580	
Body Mass Index (BMI)			
Underweight	18.38	124989	
Normal	57.74	420415	
Overweight	23.88	153957	

Table 1: Background characteristics of women aged 15-49 years surveyed.

HYPERTENSION AMONG WOMEN IN INDIA

Background Characteristics	Women surveyed		
	Per cent	Number	
Diabetes			
No	97.33	707116	
Yes	2.67	16999	
Uses of Cooking Fuel			
Clean Fuel	55.88	367560	
Polluting Fuel	44.12	356555	
Self-Smoking			
No	96.04	678443	
Yes	3.96	45672	
Exposure to Passive Smoking			
No	52.27	362507	
Yes	47.73	361608	
Uses of Tobacco			
No	97.26	690703	
Yes	2.74	33412	
Total	100	724115	

Source: Authors' calculations.

Remarks: The sample size for BMI was 699361 as BMI for 24754 women was missing.

The prevalence of hypertension among women aged 15-49 years has been found to be different in different regions of the country. The central region of the country comprising of the states of Bihar, Chhattisgarh, Jharkhand, Madhya Pradesh, and Uttar Pradesh has relatively the highest prevalence of hypertension whereas the prevalence was relatively the lowest in the west region. The prevalence of hypertension in women has also been found to be relatively high in the north and south regions compared to east and northeast regions of the country.

Table 2: Prevalence of hypertension by background characteristics of women.

Background Characteristics	Hypertension		
	Prevalence	χ^2	Ν
	(per cent)		
Age		350.14 (p<0.05)	
15-29	7.19		359559
30-44	18.25		280497
45-49	29.70		84059
Education		624.20 (p<0.05)	
No Education	19.43		167304
Primary	17.41		84983
Secondary	12.13		370012
Higher	10.40		101816
Religion		717.27 (p<0.05)	
Hindu	13.92		546007
Muslim	14.14		90729
Others	17.07		87379

Background Characteristics	Hypertension				
_	Prevalence	χ^2	Ν		
	(per cent)				
Caste		39.10 (p<0.05)			
Scheduled Caste	13.93		139957		
Scheduled Tribe	13.20		135239		
OBC*	14.20		276881		
Others	14.46		172038		
Wealth Index		672.07 (p<0.05)			
Poorest	12.84		149844		
Poorer	13.46		160340		
Middle	14.41		151505		
Richer	14.70		139607		
Richest	15.03		122819		
Regions		671.15 (p<0.05)			
North	14.96		147615		
Central	15.04		184017		
East	13.47		104342		
Northeast	13.86		103433		
West	11.65		69811		
South	14.37		114897		
Residence		190.65 (p<0.05)			
Urban	14.48	(1)	179535		
Rural	13.94		544580		
BMI***		235.74 (p<0.05)			
Underweight	8.21	(1)	124989		
Normal	12.42		420415		
Overweight	25.24		153957		
Diabetes		725.25 (p<0.05)			
No	13.46	(*)	707116		
Yes	38.04		16999		
Uses of Cooking Fuel		792.63 (p<0.05)			
Clean Fuel	14.96	(*)	367560		
Polluting Fuel	13.04		356555		
Self-Smoking		580.09 (p<0.05)			
No	13.92	()	678443		
Yes	18.88		45672		
Exposure to Passive Smoking		137.14 (p<0.05)			
No	14.43	()	362507		
Yes	13.77		361608		
Uses of Tobacco		388.85 (p<0.05)			
No	13.98	(*)	690703		
Yes	19.00		33412		
Total	14.11		724115		

Source: Authors' calculations

Figure 1 illustrates the spatial clustering of the prevalence of hypertension among women across states and union territories. The overall prevalence of hypertension in the country is estimated to be around 14.1 per cent. The prevalence of hypertension among women, however, is relatively the highest in Sikkim (25.2 per cent) followed by Punjab (23.6 per cent) and Arunachal Pradesh (22.2 per cent) but lowest in Dadra and Nagar Haveli (8.7 per cent).



Figure 1: Prevalence (per cent) of hypertension in states and Union Territories of India, 2019-2021. Source: Authors

Figure 2 depicts the variation in the prevalence of hypertension in women across districts. High prevalence of hypertension is found in districts of northeastern, and northern region, whereas low prevalence is found in districts of western and central regions. A few districts from the southern region also have high prevalence of hypertension. Moderate and high prevalence may also be observed in some of the districts of western and eastern regions. The prevalence of hypertension is found to be the highest (30.9 per cent) in North district of Sikkim but the lowest (2.4 per cent) in Shupiyan district of Jammu & Kashmir.



Figure 2: Variation in the prevalence of hypertension in women across districts of India, 2019-2021.

Source: Authors

Results of the logistic regression analysis are presented in table 3 in terms of unadjusted odds ratios (UOR) and adjusted odds ratios (AOR). We have presented both UOR and AOR to understand the independent effect of the explanatory variables on the prevalence of hypertension. The UOR shows the crude association between the cofounder and the outcome variable without controlling other factors, while AOR reflects the associations after controlling the potential cofounders to assess true effects because it prevents the distortion resulting from the external factors.

Background	Hypertension			
characteristics	UOR [CI: 95%]	ʻp'	AOR [CI: 95%]	ʻp'
Age				
15-29®	1		1	
30-44	2.883 [2.838-2.929]	0.000	2.270 [2.230-2.310]	0.000
45-49	5.456 [5.351-5.563]	0.000	4.003 [3.914-4.094]	0.000
Education				
No Education	2.077 [2.031-2.125]	0.000	1.330 [1.293-1.368]	0.000
Primary	1.816 [1.769-1.864]	0.000	1.299 [1.261-1.339]	0.000
Secondary	1.188 [1.163-1.214]	0.000	1.139 [1.112-1.166]	0.000
Higher secondary®	1		1	
Religion				
Hindu®	1		1	
Muslim	1.018 [0.998-1.038]	0.072	1.053 [1.031-1.076]	0.000
Others	1.273 [1.238-1.308]	0.000	1.267 [1.229-1.306]	0.000
Caste				
Scheduled Caste	0.957 [0.939-0.976]	0.009	1.016 [0.994-1.038]	0.163
Scheduled Tribe	0.899 [0.876-0.922]	0.000	1.099 [1.068-1.131]	0.000
OBC*	0.978 [0.963-0.995]	0.000	1.024 [1.005-1.042]	0.012
General®	1		1	
Wealth Index				
Poorest®	1		1	
Poorer	1.056 [1.033-1.079]	0.000	1.046 [1.022-1.071]	0.000
Middle	1.142 [1.118-1.167]	0.000	1.083 [1.055-1.111]	0.000
Richer	1.169 [1.145-1.195]	0.000	1.079 [1.048-1.110]	0.000
Richest	1.201 [1.175-1.123]	0.000	1.066 [1.031-1.102]	0.000
Residence				
Urban®	1		1	
Rural	0.956 [0.942-0.969]	0.000	1.034 [1.016-1.052]	0.000
BMI**				
Underweight®	1		1	
Normal	1.585 [1.550-1.620]	0.000	1.274 [1.245-1.304]	0.000
Overweight	3.774 [3.688-3.862]	0.000	2.479 [2.418-2.541]	0.000
Diabetes	4		4	
NO®		0.000	1	0.000
res	3.948 [3.832-4.068]	0.000	2.257 [2.187-2.330]	0.000

Table 3: Binary logistic regression model showing associations and determinants of hypertension among women in India.

DOLUI ET AL; IJPD 4(2): 183-199

Background	Hypertension			
characteristics	UOR [CI: 95%]	ʻp'	AOR [CI: 95%]	ʻp'
Uses of Cooking Fuel				
Clean Fuel®	1		1	
Polluting Fuel	0.852 [0.841-0.864]	0.000	0.916 [0.899-0.934]	0.000
Self-Smoking				
No®	1		1	
Yes	1.440 [1.397-1.484]	0.000	1.144 [1.080-1.211]	0.000
Exposure to Passive Sme	oking			
No®	1		1	
Yes	0.947 [0.935-0.959]	0.000	0.929 [0.916-0.942]	0.000
Uses of Tobacco				
No®	1		1	
Yes	1.443 [1.392-1.497]	0.000	0.991 [0.926-1.06]	0.000
Pseudo R ²			0.084	
Chi-Square			48508.309	
Akaike crit. (AIC)			529704.406	
Bayesian crit. (BIC)			530025.228	
Log-likelihood			-264824.2	

Source: Authors' calculations

Note: CI - Confidence interval at 95 per cent significance level; ® - Reference category; OBC - Other Backward Classes; BMI - Body Mass Index; AIC - Akaike's information criterion; BIC - Bayesian information criterion.

It may be seen from table 3 that the odd of having hypertension is more than four times higher in women aged 45-49 years compared to women aged 15-29 years as revealed through the adjusted odds ratio. On the other hand, the odd of having hypertension is around 33 per cent higher in women having no education as compared to women having more than secondary level education after controlling other factors. On the contrary, the odd of having hypertension is found to be the highest in women with middle standard of living but the lowest in women with poor level of standard of living relative to the odd of having hypertension in women with the poorest standard of living. The difference in the odd of having hypertension in women of different standard of living, however, has not been found to be large. The table also suggests that women living in the rural areas are less likely to have hypertension as compared to women living in the urban areas. However, when other factors are controlled, rural women have relatively higher chance of having hypertension compared to urban women. Obese women definitely have higher chance of having hypertension compared to normal women. Similarly, women having diabetes have higher chance of having hypertension compared to non-diabetic women. The chance of having hypertension is definitely higher in smoking women as compared to non-smoking women but the chance of having hypertension is relatively lower in passive smokers compared to women who are not passive smokers. On the basis of UOR, Scheduled Tribes women residing in the rural areas are less likely to have hypertension as compared to women of General Class but, on the basis of AOR, they are more likely to have hypertension compared to General Class women of the rural areas. However, the AOR is found to be statistically insignificant,

Discussion

Hypertension is a prominent risk factor for cardiovascular diseases and frequently contributes to multimorbidity (Dolui et al, 2023; Mohanty et al, 2021). The prevalence of hypertension has been found to be the highest among all morbidities (Dolui et al, 2023). The present analysis has unveiled several noteworthy findings. Our analysis confirms that prevalence of hypertension increases with age. It is well-known that the prevalence of hypertension was significantly higher among women of older ages compared to women of younger ages (Pires et al, 2013). Our finding aligns with similar observations in India, where the prevalence of hypertension is found to be higher among adults in the age group 45 years and above (Geldsetzer et al, 2018; Meshram et al, 2022; Mohanty et al, 2021). This association can be attributed to the progressive increase in mean systolic blood pressure throughout adulthood, with a steeper rise of blood pressure observed in women compared to men (Kearney et al, no date; Ong et al, 2008; Pimenta, 2011; Roger et al, 2011; Westheim et al, 2001).

We have also found that educational level of women is negatively associated with the prevalence of hypertension. This finding aligns with a previous study, which has also demonstrated that individuals with below-average educational level have higher risk of hypertension (Sun et al, 2022). It has also been observed that, for each additional year of education, there is a corresponding decrease in the systolic blood pressure, which reduces the likelihood of hypertension (Liu et al. 2011). The risk of hypertension is also found to be low in Scheduled Tribes and Other Backward Classes (OBC), exhibited lower awareness and were more prone to hypertension. Additionally, Women from Scheduled tribe categories are likely to be part of the causes of the high prevalence of hypertension, as suggested by previous studies (Anchala et al, 2014; Chakma et al, 2017; Gupta, 2004; Laxmaiah et al, 2015; Rizwan et al, 2014). Similarly, some studies revealed that women from tribal backgrounds with lower educational attainment face a higher risk of hypertension (Gupta et al. 1994: Hajjar et al, 2001; Kopp, 2005; Laxmaiah et al, 2015; Stamler et al, 1996, 2002). The present study also confirms that the risk of hypertension is directly related to the standard of living - the higher the standard of living the higher the prevalence of hypertension as observed in other studies (Mohanty et al, 2021).

Likewise, our study has revealed that women in the urban areas have higher risk of hypertension compared to women in the rural areas. An earlier study (Gupta et al, 1995) has also found higher risk of hypertension in urban women. However, the adjusted odds ratio suggests that the risk of hypertension is higher in rural women than in urban women which is also supported by various studies (Bisquera et al, 2022; Boro and Banerjee, 2022; Busingye et al, 2017; Chauhan et al, 2021; Gupta, 2004). The risk of hypertension has also been found to be directly associated with the obesity in women as revealed through BMI. A similar study has also reported that obese individuals with hypertension are at a greater risk of developing coronary heart disease than the non-obese one (Chiang et al, 1969). On the contrary, some studies have highlighted that low BMI can also be a contributing risk factor for hypertension, particularly among people in rural areas who may have low BMI because of chronic micronutrient deficiencies (Chakma et al, 2017; Dolui, et al, 2023; Goldbourt et al, 1987; Hu et al, 2000; Tesfaye et al, 2007).

The prevalence of hypertension among diabetic women is notably higher and regressively associated in various aspects when compared to other groups. These findings align with those from earlier studies (Chauhan et al, 2021; Geldsetzer et al, 2018). Additionally, a significant risk factor for hypertension among women is self-smoking. Earlier studies have shown that maternal smoking or smoking during pregnancy leads to the risk of hypertension (Liang et al, 2022; Seal et al, 2013). However, the present analysis suggests passive smoking does not appear to be a risk factor in hypertension.

The present study has certain limitations. First, it is based on the cross-sectional data, which may not capture the long-term trend in the prevalence of hypertension or causality. Second, the study is limited to women aged 15-49 years only, which may not fully represent the entire women population. The National Family Health Survey has also collected data on the blood pressure for men covered under the survey and there are several studies that have analysed the prevalence of hypertension and its determinants in both men and women using the NFHS data. In the present analysis, however, we have focused, particularly, on women aged 16-49 years to find out how selected demographic and socio-economic characteristics of these women impact upon the level of blood pressure. These limitations should be considered when interpreting the findings of the present analysis and may offer opportunities for further research and data collection to address these gaps.

Conclusion

The present study draws attention to the pressing issue of hypertension among women in India and offers crucial insights into the chronic burden of women, which contributes substantially to the prevalence of cardiovascular diseases and multimorbidity. The findings emphasise that elderly women, those with lower levels of education, and belonging to the socioeconomically disadvantaged section of the community, face a higher risk of hypertension. The prevalence of hypertension may lead to other non-communicable diseases, such as diabetes, and reveals the impact of smoking on blood pressure. Recognising these vulnerability factors is of utmost importance for the development of effective public health interventions aimed at reducing the prevalence of hypertension and associated risk factors in women. The government of India has launched the India Hypertension Control Initiative (IHCI) in 2017 which aims aims to accelerate progress towards the Non-Communicable Diseases (NCD) target set by the Government of India by supplementing and intensifying evidence-based strategies to strengthen the building blocks of hypertension management and control. The Government of India aims to reduce premature mortality due to non-communicable diseases (NCDs) by 25 per cent by 2025. One of the nine voluntary targets is to reduce the prevalence of high blood pressure by 25 per cent by 2025. The present study suggests that target-specific interventions based on education, wealth and age such as awareness campaigning for quitting smoking substances, healthy dietary patterns, physical activities, community diagnosis and screening and easy access to low-cost or free antihypertensive medication is necessary to manage hypertension in women.

References

- Abarca-Gómez L, Abdeen ZA, Hamid ZA, Abu-Rmeileh NM, Acosta-Cazares B, Acuin C, Adams RJ, Aekplakorn W, Afsana K, Aguilar-Salinas CA, Agyemang C, Ahmadvand A, Ahrens W, Ajlouni K, Akhtaeva N, Al-Hazzaa HM, Al-Othman AR, Al-Raddadi R, Al Buhairan F, ... Ezzati M (2017) Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128-9 million children, adolescents, and adults. *The Lancet* 390(10113), 2627–2642.
- Anchala R, Kannuri NK, Pant H, Khan H, Franco OH, Di Angelantonio E, Prabhakaran D (2014) Hypertension in India: a systematic review and meta-analysis of prevalence, awareness, and control of hypertension. *Journal of Hypertension* 32(6): 1170-1177.
- Bhimarasetty MD, Pamarthi K, Prasad Kandipudi KL, Padmasri Y, Nagaraja SB, Khanna P, Goel S (2022) Hypertension among women in reproductive age in India: Can we predict the risk? An analysis from National Family Health Survey (2015-2016). *Journal of Family Medicine and Primary Care* 11(9), 5857–5864.
- Bisquera A, Turner EB, Ledwaba-Chapman L, Dunbar-Rees R, Hafezparast N, Gulliford M, Durbaba S, Soley-Bori M, Fox-Rushby J, Dodhia H, Ashworth M, Wang Y (2022) Inequalities in developing multimorbidity over time: A population-based cohort study from an urban, multi-ethnic borough in the United Kingdom. *The Lancet Regional Health Europe* 12: 1-12.
- Borah PK, Mahanta J, Kalita HC, Bhattacharjee CK (2020) Distribution of hypertension subtypes in a hill tribe of Mizoram, Northeast India. *Indian Heart Journal* 72(5): 398–402.
- Boro B, Banerjee S (2022) Decomposing the rural-urban gap in the prevalence of undiagnosed, untreated and under-treated hypertension among older adults in India *BMC Public Health* 22(1): 1-16.
- Busingye D, Arabshahi S, Evans RG, Srikanth VK, Kartik K, Kalyanram K, Riddell MA, Zhu X, Suresh O, Thrift AG (2017) Factors associated with awareness, treatment and control of hypertension in a disadvantaged rural Indian population. *Journal of Human Hypertension* 31(5), 347–353.
- Chakma T, Kavishwar A, Sharma RK, Rao PV (2017) High prevalence of hypertension and its selected risk factors among adult tribal population in Central India. *Pathogens and Global Health* 111(7): 343-350.
- Chauhan S, Gupte SS, Kumar S, Patel R (2021) Urban-rural differential in diabetes and hypertension among elderly in India: A study of prevalence, factors, and treatment-seeking. *Diabetes & Metabolic Syndrome* 15(4): 1-8.

- Chiang BN, Perlman LV, Epstein FH (1969) Overweight and hypertension. *Circulation* 39(3): 403–421.
- Daniel RA, Haldar P, Kant S, Krishnan A, Kumar R, Gupta SK (2022) Prevalence and factors associated with hypertension among community-dwelling adolescents in a rural area of Haryana, India, using AAP 2017 and NHBPEP criteria. *Indian Journal of Pediatrics* 89(11): 1099–1106.
- Dolui M, Sarkar S, Ghosh P, Hossain M (2023) Dietary diversity and association with noncommunicable diseases (NCDs) among adult men (15–54 years): A cross-sectional study using National Family and Health Survey, India. *PLOS Global Public Health* 3(4): 1-17.
- Dolui M, Sarkar S, Hossain M, Manna H (2023) Demographic and socioeconomic correlates of multimorbidity due to non-communicable diseases among adult men in India: Evidence from the nationally representative survey (NFHS-5). *Clinical Epidemiology and Global Health* 23(101376): 1–9.
- Ezzati M, Lopez AD, Rodgers A, Vander Hoorn S, Murray CJ (2002) Selected major risk factors and global and regional burden of disease *The Lancet* 360(9343): 1347–1360.
- Geevar Z, Krishnan MN, Venugopal K, Sanjay G, Harikrishnan S, Mohanan PP, Mini GK, Thankappan KR (2022) Prevalence, awareness, treatment, and control of hypertension in young adults (20–39 Years) in Kerala, South India. *Frontiers in Cardiovascular Medicine*, 9: 1-11.
- Geldsetzer P, Manne-Goehler J, Theilmann M, Davies JI, Awasthi A, Vollmer S, Jaacks LM, Bärnighausen T, Atun R (2018) Diabetes and hypertension in India: a nationally representative study of 13 million adults. *JAMA Internal Medicine* 178(3): 363–372.
- Goldbourt U, Holtzman E, Cohen-Mandelzweig L, Neufeld HN (1987) Enhanced risk of coronary heart disease mortality in lean hypertensive men. *Hypertension* 10(1): 22–28.
- Gupta R (2004) Trends in hypertension epidemiology in India. *Journal of Human Hypertension* 18(2): 73–78.
- Gupta R, Gupta VP, Ahluwalia NS (1994) Educational status, coronary heart disease, and coronary risk factor prevalence in a rural population of India. *BMJ*: *British Medical Journal* 309(6965): 1332-1336.
- Gupta R, Gupta S, Gupta VP, Prakash H (1995) Prevalence and determinants of hypertension in the urban population of Jaipur in western India. *Journal of Hypertension* 13(10): 1193-1200.

- Hajjar IM, Grim CE, George V, Kotchen, TA (2001) Impact of diet on blood pressure and agerelated changes in blood pressure in the US population: analysis of NHANES III. *Archives of Internal Medicine*, 161(4), 589–593.
- Hu FB, Wang B, Chen C, Jin Y, Yang J, Stampfer MJ, Xu X (2000) Body mass index and cardiovascular risk factors in a rural Chinese population. *American Journal of Epidemiology* 151(1): 88–97.
- Government of India (2021) *National Family Health Survey (NFHS-5) 2019-21. India*. New Delhi, Ministry of Health and Family Welfare, International Institute for Population Sciences.
- Jonas JB, Nangia V, Matin A, Joshi PP, Ughade SN (2010) Prevalence, awareness, control, and associations of arterial hypertension in a rural central India population: The central India eye and medical study. *American Journal of Hypertension* 23(4): 347–350.
- Kearney PM, Whelton M, Reynolds K, Muntner P, Whelton PK, He J (2005) Global burden of hypertension: analysis of worldwide data. *The Lancet*, 365(9455), 217–223.
- Kopp W (2005) Pathogenesis and etiology of essential hypertension: role of dietary carbohydrate. *Medical Hypotheses* 64(4): 782–787.
- Laxmaiah A, Meshram II, Arlappa N, Balakrishna N, Mallikharjuna Rao K, Reddy CG, Ravindranath M, Kumar S, Kumar H, Brahmam GNV (2015) Socio-economic & demographic determinants of hypertension & knowledge, practices & risk behaviour of tribals in India. *Indian Journal of Medical Research* 141(5): 697–708.
- Liang J, Fu Z, Liu Q, Shen Y, Zhang X, Weng Z, Xu J, Li W, Xu C, Zhou Y, Gu A (2022) Interactions among maternal smoking, breastfeeding, and offspring genetic factors on the risk of adult-onset hypertension. *BMC Medicine* 20(1): 1–11.
- Liu SY, Buka SL, Linkletter CD, Kawachi I, Kubzansky L, Loucks EB (2011) The association between blood pressure and years of schooling versus educational credentials: Test of the sheepskin effect. *Annals of Epidemiology* 21(2): 128: 138.
- Loke WH, Ching SM (2022) Prevalence and factors associated with psychological distress among adult patients with hypertension in a primary care clinic: A cross-sectional study. *Malaysian Family Physician* 17(2): 89–98.
- Meshram I, Boiroju NK, Longvah T (2022) Prevalence of overweight/obesity, hypertension and its associated factors among women from Northeast India. *Indian Heart Journal* 74(1): 56–62.
- Meshram II, Vishnu Vardhana Rao M, Sudershan Rao V, Laxmaiah A, Polasa K (2016) Regional variation in the prevalence of overweight/obesity, hypertension and diabetes and their correlates among the adult rural population in India. *British Journal of Nutrition* 115(7): 1265–1272.

- Mishra VK, Srivastava S, Muhammad T, Murthy PV (2022) Relationship between tobacco use, alcohol consumption and non-communicable diseases among women in India: evidence from National Family Health Survey-2015-16. *BMC Public Health* 22(1): 1-12.
- Mohanty SK, Pedgaonkar SP, Upadhyay AK, Kämpfen F, Shekhar P, Mishra RS, Maurer J, O'Donnell O (2021) Awareness, treatment, and control of hypertension in adults aged 45 years and over and their spouses in India: A nationally representative crosssectional study/ *PLOS Medicine* 18(8): 1-20.
- Government of India (2022) *National Multisectoral Action Plan (NMAP) for Prevention and Control of Common NCDs (2017-22)*. New Delhi, Ministry of Health and Family Welfare.
- Ong KL, Tso AWK, Lam KSL, Cheung BMY (2008) Gender difference in blood pressure control and cardiovascular risk factors in Americans with diagnosed hypertension. *Hypertension* 51(4 Part 2 Suppl): 1142–1148.
- Pimenta E (2011) Hypertension in women. Hypertension Research 35(2): 148-152.
- Pires JE, Sebastião YV, Langa AJ, Nery SV (2013) Hypertension in Northern Angola: Prevalence, associated factors, awareness, treatment and control. *BMC Public Health* 13(1): 1–10.
- Rizwan SA, Kumar R, Singh AK, Kusuma YS, Yadav K, Pandav CS (2014) Prevalence of hypertension in Indian Tribes: a systematic review and meta-analysis of observational studies. *PLoS ONE* 9(5): 1-10.
- Roger VL, Go AS, Lloyd-Jones DM, Adams RJ, Berry JD, Brown TM, Carnethon MR, Dai S, de Simone G, Ford ES, Fox CS, Fullerton HJ, Gillespie C, Greenlund KJ, Hailpern SM, Heit JA, Michael HP, Howard VJ, Kissela BM, ... Wylie-Rosett J (2011) Heart disease and stroke statistics - 2011 update: a report from the American Heart Association. *American Heart Association* 123(4): 18-209.
- Saha I, Paul B, Dasgupta A (2008) Prevalence of hypertension and variation of blood pressure with age among adolescents in Chetla, India. *Tanzania Journal of Health Research* 10(2): 108-111.
- Seal N, Krakower G, Seal J (2013) Maternal smoking during pregnancy and metabolic syndrome in their children. *The Journal for Nurse Practitioners* 9(10): 695–705.
- Stamler J, Caggiula A, Grandits GA, Kjelsberg M, Cutler JA (1996) Relationship to blood pressure of combinations of dietary macronutrients. Findings of the Multiple Risk Factor Intervention Trial (MRFIT). *Circulation* 94(10): 2417–2423.
- Stamler J, Liu K, Ruth KJ, Pryer J, Greenland P (2002) Eight-year blood pressure change in middle-aged men: relationship to multiple nutrients. *Hypertension* 39(5): 1000–1006.

- Sun K, Lin D, Li M, Mu Y, Zhao J, Liu C, Bi Y, Chen L, Shi L, Li Q, Yang T, Wan Q, Wu S, Wang G, Luo Z, Qin Y, Tang X, Chen G, Huo Y, ... Yan L (2022) Association of education levels with the risk of hypertension and hypertension control: a nationwide cohort study in Chinese adults. *Journal of Epidemiology and Community Health* 76(5): 451–457.
- Tesfaye F, Nawi NG, Van Minh H, Byass P, Berhane Y, Bonita R, Wall S (2007) Association between body mass index and blood pressure across three populations in Africa and Asia. *Journal of Human Hypertension* 21(1): 28–37.
- Westheim A, Klemetsrud T, Tretli S, Stokke HP, Olsen H (2001) Blood pressure levels in treated hypertensive patients in general practice in Norway. *Blood Pressure* 10(1): 37–42.
- WHO (2015) Noncommunicable diseases: Hypertension. Geneva, World Health Organization.
- WHO (2023) *Global Report on Hypertension. The Race against a Silent Killer*. Geneva, World Health Organization.